

AMERICAN SOCIETY FOR REPRODUCTIVE MEDICINE

Infertility: An Overview

A Guide for Patients



PATIENT INFORMATION SERIES

Published by the American Society for Reproductive Medicine under the direction of the Patient Education Committee and the Publications Committee. No portion herein may be reproduced in any form without written permission. This booklet is in no way intended to replace, dictate, or fully define evaluation and treatment by a qualified physician. It is intended solely as an aid for patients seeking general information on issues in reproductive medicine.

Copyright © 2012 by the American Society for Reproductive Medicine

Infertility: An Overview

A Guide for Patients Revised 2012

A glossary of italicized words is located at the end of this booklet.

INTRODUCTION

Infertility is typically defined as the inability to achieve pregnancy after one year of unprotected intercourse. If you have been trying to conceive for a year or more, you should consider an infertility evaluation. However, if you are 35 years or older, you should begin the infertility evaluation after about six months of unprotected intercourse rather than a year, so as not to delay potentially needed treatment. If you have a reason to suspect an underlying problem, you should seek care earlier. For instance, if you have very irregular menstrual cycles (suggesting that you are not ovulating or releasing an egg), or if you or your partner has a known fertility problem, you probably should not wait an entire year before seeking treatment. If you and your partner have been unable to have a baby, you're not alone. At least one out of seven couples has trouble conceiving. During this time, it is normal to experience feelings of frustration, jealousy, anger and stress. However, once you begin to explore your medical options, you'll find that fertility treatments offer more hope for a successful pregnancy than ever before. The vast majority of patients who seek care from a fertility specialist reach their goal of becoming pregnant.

THE INITIAL VISIT

Your regular gynecologist may perform the initial fertility evaluation, or you may choose to be evaluated by an infertility specialist. It is important to find a physician who you trust and with whom you feel comfortable. Some general obstetricians/gynecologists (ob/gyns) have a special interest in the treatment of infertility. Reproductive endocrinologists are specialized ob/gyns who, after completing the full ob/gyn residency, complete three additional years of specialized training in reproductive endocrinology and infertility at an approved fellowship program. If an abnormal *semen analysis* is known or later found, the male partner should be referred to a urologist who specializes in the area of male infertility. To locate physicians in your area who specialize in female or male infertility issues, go to the ASRM patient website at:

www.ReproductiveFacts.org and click on “Find a Healthcare Professional” on the right side of the page. You can search for a health professional by name, city, and specialty.

Once you choose a qualified physician, you and your partner will undergo a series of tests. Sometimes this may require a significant amount of time and energy on your part, but the results are invaluable. Before these tests are performed, your physician will ask questions and review any records that you and your partner may have from a previous infertility evaluation. Ensuring that your physician has access to review previous medical records is critical in minimizing wasted time and money for repeated diagnostic evaluation and therapy. Both you and your partner should try to attend the first meeting since infertility is a shared experience and is best dealt with as a couple. During this visit, you’ll begin to understand the degree of commitment and cooperation that an infertility investigation requires. It is important that you communicate freely between each other as well as with your physician.

Your physician will review your history and ask you and your partner questions to help identify potential causes for your difficulty in conceiving. In a typical initial visit, the physician will ask about the frequency and regularity of your menstrual period, pelvic pain, abnormal vaginal bleeding or discharge, history of pelvic infection, and medical illnesses. Expect questions concerning prior pregnancies, miscarriages, operations, and methods of contraception. Your partner will be asked questions concerning prior genital injury, surgery, infections, drug and/or medication use, history of fathering other children, and medical illnesses. You’ll be asked how long you’ve been trying to conceive, how often you have sexual intercourse, if you use lubricants during intercourse, and if anyone in either family has birth defects. Your physician will need to know the complete sexual and reproductive histories of you and your partner, including any former relationships. Since at least 25% of infertile couples have more than one factor causing infertility, it is very important to evaluate all factors that may affect both you and your partner.

During the first visit, you may discuss the emotional stress of infertility, a subject that is often difficult for you to share with family and friends. Physicians know that the procedures and intimate questions involved in an infertility workup can be difficult. You should always feel free to make your physician aware of your concerns and frustrations and you should ask your physician questions whenever you need clarification.

A physical examination of you and your partner may follow the initial review. The occurrence and extent of the examinations will depend upon whether or not any factors impacting your fertility are found early in the evaluation.

THE PROCESS OF CONCEPTION

In order to understand the fertility tests and treatments you may be offered, it is important to understand how conception naturally occurs. First, your ovary must release (ovulate) an egg, which must be picked up by the fallopian tube. Sperm must travel through the *vagina*, into the *uterus*, and up into the fallopian tube in order to fertilize the egg. Fertilization usually takes place in the fallopian tube. Then, the fertilized egg, or embryo, travels down the fallopian tube to the uterus, where it implants in the uterine lining and develops (Figure 1). A problem that develops in any part of this process can lead to infertility.

Figure 1




Figure 1. Arrows indicate the path the sperm and egg must take to meet. The fertilized egg continues traveling through the fallopian tube to the uterus.

However, some women who ovulate normally may not have a rise in temperature, and many factors unrelated to the reproductive cycle, such as a cold or fatigue, can affect the BBT. At best, the BBT chart helps determine when and if ovulation is taking place, but only after it has happened.

To predict ovulation before it takes place, in order to schedule intercourse or insemination, for example, you may use an over-the-counter ovulation prediction kit. These urine test kits are designed to detect the surge of *luteinizing hormone (LH)* that occurs just before you ovulate. The LH surge stimulates one of the ovaries to release an egg and produce progesterone. Ovulation prediction kits usually detect the LH surge about a day or a day and a half before ovulation, giving you and your partner advance notice of an egg that will be released (ovulation). However, not all women who ovulate will have an LH surge that will be detected using these kits. You may find these kits difficult and frustrating, and false positive and false negative results do occur occasionally.

In a normal cycle, progesterone levels are highest about seven days after ovulation. Your physician may perform a blood test to measure the level of progesterone in your blood at this time. Generally, blood progesterone is tested on day 19 to 23 of a 28-day menstrual cycle. An elevated progesterone level helps to confirm ovulation and the adequacy of ovarian hormone production.

Your physician may perform a pelvic ultrasound examination to evaluate ovulation, but this is not done routinely. This examination may indicate whether your ovaries are producing *follicles*. These follicles are fluid filled sacs located just beneath the *ovary's* surface that contain the immature eggs. Ultrasound may also help to document a follicle's collapse and subsequent release of fluid, implying release of an egg.

If you are not ovulating, your physician may order special tests to determine the reason and then prescribe certain drugs to induce ovulation. Your medical history and physical exam will help determine which tests are appropriate. Both oral and injectable medications are available to help induce your ovulation.

The Tubal Factor

Because open and functional fallopian tubes are necessary for conception,

tests to determine tubal openness (patency) are important. Tubal factors, as well as factors affecting the peritoneum (lining of the pelvis and abdomen), account for about 35% of all infertility problems. A special x-ray called a *hysterosalpingogram* (HSG) (Figure 3) can be performed to evaluate the fallopian tubes and uterus (see ASRM patient factsheet titled *Hysterosalpingogram*). During an HSG, a special fluid (dye) is injected through your *cervix*, fills your uterus, and travels into your fallopian tubes. If the fluid spills out the ends of the tubes, they are open. If the fluid does not spill out the ends, then the tubes are blocked. If the HSG shows blocked fallopian tubes, your physician may perform a laparoscopy to assess the degree of tubal damage.

Figure 3




Figure 3. Hysterosalpingogram, a procedure to determine if the fallopian tubes are open or blocked.

If the tubes are found to be blocked, scarred, or damaged, surgery can sometimes correct the problem. But surgery does not guarantee that the tube, even if opened up or cleared of scar tissue, will function properly. Although some tubal problems are correctable by surgery, women with severely damaged tubes are so unlikely to become pregnant that *in vitro fertilization* (IVF) offers them the best hope for a successful pregnancy. Because very badly damaged tubes may fill with fluid (*hydrosalpinx*) and lower IVF success rates, your physician may recommend removal of the damaged tubes prior to IVF. More information on tubal disease can

be found in the ASRM Patient Information booklet titled *Tubal Factor Infertility*.

The Male Factor

In approximately 40% of infertile couples, the male partner is either the sole or a contributing cause of infertility. Therefore, a semen analysis is important in the initial evaluation. To prepare for a semen analysis, your partner will be asked to abstain from ejaculating for at least 48 hours. He then will collect a semen sample in a cup by masturbating at home or in the physician's office. In some instances, a semen sample may be obtained during sexual intercourse using a special condom provided by the physician. The semen specimen is examined under a microscope to determine the volume (amount), motility (movement), and morphology (appearance and shape) of the sperm. In general, two or three semen analyses may be recommended over two to six months, since sperm quality can vary over time. Other hormonal and genetic tests may be recommended as well depending on the type and severity of abnormalities found.

If your partner's semen analysis reveals abnormalities, he may need to consult a clinician who specializes in male infertility. Treatment for male factor infertility may include antibiotic therapy for infection, surgical correction of *varicocele* (dilated or varicose veins in the scrotum) or duct obstruction, or medications to improve sperm production. In some men, surgery to obtain sperm from the testis can be performed. In some cases, no obvious cause of poor sperm quality can be found. *Intrauterine insemination (IUI)* or IVF may then be recommended. Direct injection of a single sperm into an egg (*intracytoplasmic sperm injection [ICSI]*) may be recommended as a part of the IVF process. For more information, refer to the ASRM Patient Information booklets titled *Male Infertility* and *Vasectomy Reversal*. If no sperm are present on semen analysis or not found at surgical extraction, your physician may discuss using a sperm donor. Insemination with donor sperm may also be considered if IUI is not successful or if you and your partner do not choose to undergo IVF.

The Age Factor

Delaying pregnancy is a common choice for women in today's society. The number of women in their late 30s and 40s attempting pregnancy and having babies has increased in recent years. If you've chosen to delay pregnancy, due to college or career for example, you may not realize that your fertility begins to decline significantly in your mid 30s and accelerates

in your late 30s. Some women even begin to experience a decline in their fertility in their late 20s and early 30s.

Fertility declines with age because fewer eggs remain in your ovaries, and the quality of the eggs remaining is lower than when you were younger. Blood tests are now available to determine your *ovarian reserve*, a term which reflects your age-related fertility potential. In the simplest of these tests, the hormones *follicle-stimulating hormone (FSH)* and *estradiol* are tested in your blood on the second, third, or fourth day of your menstrual period. An elevated FSH level indicates that your chances for pregnancy may be lower than routinely expected for your age, especially if you are age 35 or older. In addition, an AMH level (*anti-müllerian hormone level*) may also be ordered to provide additional information about your ovarian reserve. A lower AMH level indicates decreased ovarian reserve. Abnormally high FSH or low AMH levels do not mean that you have no chance of successful conception. However, they may indicate that success rates may be lower, that more aggressive treatment may be warranted, and/or that higher medication doses may be needed. Another commonly used option to detect ovarian reserve is the use of transvaginal ultrasound to determine *antral follicle count (AFC)*, which is when each follicle in both ovaries are counted. An AFC is performed during the first 3-4 days of the menstrual cycle.

Older women tend to have a lower response to fertility medications and a higher miscarriage rate than younger women. The chance of having a chromosomally abnormal embryo, such as one with Down syndrome, also increases with age. Because of the marked effect of age on pregnancy and birth rates, it is common for older couples to begin fertility treatment sooner and, in some cases, to consider more aggressive treatment than younger couples.

Possible treatments for age-related infertility in women include fertility drugs plus IUI or IVF. In cases where the treatments fail or are predicted to have a low chance of success, *egg donation* is an option. Egg donation has a high chance of success, regardless of your FSH level. More recently, *embryo donation* has also become a viable option for many couples. For couples who have not succeeded with fertility treatments or who choose to forgo treatment, adoption is an important option. For more information about these options, please consult the ASRM Patient Information booklets titled *Age and Fertility* and *Adoption*.

The Cervical/Uterine Factor

Conditions within the cervix, which is the lower part of the uterus, may impact your fertility, but they are rarely the sole cause of infertility. It is important for your physician to know if you have had prior biopsies such as a cone biopsy, surgery, “freezing” and/or laser treatment of the cervix (LEEP), abnormal pap smears, or if your mother took DES (diethylstilbestrol) while she was pregnant. In order to determine if there is a problem with your cervix. Cervical problems are generally treated with antibiotics, hormones, or by IUI.

The HSG test, often used to investigate the *fallopian tubes*, can also reveal defects inside the uterine cavity, which is the hollow space inside your uterus where an embryo would implant and develop. An HSG is typically done after your period stops and before *ovulation*. Possible uterine abnormalities that may be identified include scar tissue, polyps (bunched-up pieces of the endometrial lining), *fibroids*, or an abnormally-shaped uterine cavity. Problems within your uterus may interfere with implantation of the embryo or may increase the incidence of miscarriage. *Hysteroscopy*, a minor and minimally invasive, incision-free surgical procedure, may be required to further evaluate and possibly correct uterine structural problems.

Uterine abnormalities can also be seen with a saline sonohysterogram (SHG). This test may also be called a sonohysterogram or water ultrasound. An SHG is a pelvic ultrasound performed while sterile fluid is injected through your cervix to outline the uterine cavity. Unlike the HSG, the SHG allows visualization of the wall of the uterus as well as the cavity at the same time, a difference which may be helpful in many cases.

The Peritoneal Factor

Peritoneal factor infertility refers to abnormalities involving the *peritoneum* (lining of the surfaces of your internal organs) such as scar tissue (*adhesions*) or *endometriosis* (Figure 4). Endometriosis is a condition where tissue that normally lines the uterus begins to grow outside the uterus. This tissue may grow on any structure within the pelvis including the ovaries and is found in about 35% of infertile women who have no other diagnosable infertility problem.

Figure 4




Figure 3. Endometriosis and adhesions

Endometriosis is found more commonly in women with infertility, pelvic pain, and painful intercourse. Endometriosis may affect the function of the ovaries, your ovarian reserve, the function of the fallopian tubes, as well as implantation. For further information concerning this problem, see the ASRM Patient Information booklet titled *Endometriosis*.

Laparoscopy is a same-day, minimally invasive surgical procedure that is performed to diagnose and treat adhesions or endometriosis. Laparoscopy is usually performed under general anesthesia, often in a hospital's same-day surgery unit. During the procedure, a narrow, lighted telescopic instrument called a *laparoscope* is inserted through a small incision within or just below your navel (Figure 5). The physician then looks through the laparoscope directly into your abdominal cavity and inspects the ovaries, fallopian tubes, and uterus.

A dye may be injected through the cervix to determine if the passage to the uterus and tubes is open. One or more additional smaller incisions may be made above your pubic area in order for the physician to insert additional instruments to better examine your pelvic organs and treat any disease if found.

A variety of specialized surgical instruments, including robotic-controlled instruments and lasers, may now be used laparoscopically to treat such conditions as endometriosis, adhesions, and ovarian cysts such as endometriomas (fluid-filled cysts on the ovaries). In many cases, laparotomy, which involves a larger abdominal incision than laparoscopy and an approximately six-week recovery, is no longer necessary since many problems can now be treated laparoscopically.

Figure 5




Figure 5. Laparoscopy

Although it is reasonable and sometimes advisable to perform laparoscopy to identify and treat pelvic disease, this procedure is not routinely performed in every infertility evaluation. Laparoscopy is expensive and involves some small surgical risk. You and your physician should discuss whether or not laparoscopy is recommended in your particular case. Sometimes your physician may not recommend a laparoscopy initially, but may instead recommend it at a later time depending on your response to treatment.

Unexplained Infertility

In approximately 10% of couples trying to conceive, all of the above tests are normal and there is no easily identifiable cause for infertility. In a much higher percentage of couples, only minor abnormalities are

found that should not be severe enough to result in infertility. In these cases, the infertility is referred to as “unexplained”. Couples with unexplained infertility may have problems with egg quality, fertilization, genetics, tubal function, or sperm function that are difficult to diagnose and/or treat. Fertility drugs and IUI have been used in couples with unexplained infertility with limited success. If no pregnancy occurs within three to six treatment cycles, IVF may be recommended and has been shown to be the most effective treatment for unexplained infertility.

GENETIC ABNORMALITIES

Some men and woman may carry genetic abnormalities that make it more difficult to become pregnant and more likely that a pregnancy ends in miscarriage. One example is a translocation, or a rearrangement of genetic material. This may be tested for, in appropriate circumstances, by blood testing of the couple. Some couples may even carry a known genetic illness and wish to avoid passing this illness on to a child. In special circumstances, a more advanced fertility treatment known as *pre-implantation genetic diagnosis (PGD)* may be offered in conjunction with IVF. PGD allows the couple to find out genetic information about embryos prior to implanting those embryos in a woman’s body. This treatment is only helpful in certain couples and requires specific genetic counseling. Your physician may discuss this form of treatment depending on your individual situation.

CHANCES OF SUCCESS WITH IVF

Each couple has a unique set of circumstances, and the chances of treatment success vary widely. The success of IVF depends on many factors, especially your age and the quality of your embryos. For more information on IVF, refer to the ASRM Patient Information booklet titled *Assisted Reproductive Technologies*. National and regional assisted reproductive technologies (ART) success rates are available on the Internet at: www.cdc.gov and www.sart.org.

PSYCHOLOGICAL IMPLICATIONS

Infertility is a medical condition that has many emotional aspects. Feelings such as anger, sadness, guilt, and anxiety are common and may affect your self-esteem and self-image. You may find it difficult to share your feelings with family and friends, which can lead to isolation. It is important to know that these feelings are normal responses to infertility and are experienced by many couples. Although a physician will describe various treatments and realistic odds of success with treatments, you must decide how far you will go in your attempts to conceive. Coming to a joint decision with

your partner about goals and acceptable therapies is important. Setting endpoints for therapy may also be advisable. For more information, visit the ASRM patient education website at www.ReproductiveFacts.org and the ASRM Mental Health Professionals Group website at www.asrm.org/MHPG_Patient_Resources. You may also go to the ASRM website at www.asrm.org or www.ReproductiveFacts.org and click on “Find a Healthcare Professional” to locate counselors in your area who specialize in infertility issues.

SUMMARY

Sometimes the factors affecting your fertility are easy to detect and treat, but in many cases a specific reason for infertility may be difficult to identify. After a full evaluation, your physician can give you a reasonable idea of your chances of achieving pregnancy with various treatment options. The choice of which treatment to pursue, if any, is strictly a personal one. Side effects, costs, the risk of multiple births, which is increased with some infertility treatments, and expected success rates are important factors to consider when choosing a treatment plan. It is helpful when beginning fertility treatment to develop a long-term plan with your physician so that you will have an idea of how long to pursue a particular treatment before moving on to more aggressive therapy or stopping treatment. Since your initial treatment may provide additional answers as to the cause of your infertility, it is very important to continually re-evaluate your plan by discussing treatment results with your physician. It may be necessary at times to temporarily cease therapy and begin again at a later date. Thanks to the many options existing today, including advanced reproductive technologies and adoption, most infertile couples will be able to experience the joy of parenthood. The sooner you meet with a specialist, the sooner you will find answers and be able to realize your dream of having a child.

GLOSSARY

Adhesions. Bands of fibrous scar tissue that may bind the pelvic organs and/or loops of bowel together. Adhesions can result from infections, endometriosis, or previous surgeries.

Anti-Müllerian hormone (AMH). A hormone measurement that reflects your ovaries' capability for producing eggs.

Antral follicle count (AFC). A transvaginal ultrasound machine is used to determine the number of pre-stimulated follicles in the ovary during the first 3-4 days of the menstrual cycle.

Assisted Reproductive Technologies (ART). All treatments that include laboratory handling of eggs, sperm, and/or embryos. The most common examples of ART are in vitro fertilization (IVF), intracytoplasmic sperm injection (ICSI), pre-implantation genetic diagnosis (PGD), frozen embryo transfer (FET), donor egg, and donor embryo transfer.

Basal body temperature (BBT). The body temperature at rest. It is taken orally each morning immediately upon awakening and recorded on a BBT chart. The readings are studied to help identify ovulation, which usually occurs at approximately the same time as the rise in BBT.

Cervix. The narrow, lower part of the uterus that opens into the vagina. The cervical canal runs through the cervix and connects the vagina with the uterine cavity. The cervix produces mucus through which sperm must swim before entering the uterine cavity and then the fallopian tubes.

Corpus luteum. A mature follicle that has collapsed after releasing its egg at ovulation. The corpus luteum secretes progesterone and estrogen during the second half of a normal menstrual cycle. The secreted progesterone prepares the lining of the uterus (endometrium) to support a pregnancy.

Egg Donation. The process of fertilizing eggs from a donor with the male partner's sperm in a laboratory dish and transferring the resulting embryos to the female partner's uterus. The female partner will not be biologically related to the child, although she will be the birth mother on record. The male partner will be biologically related to the child.

Embryo Donation. The process of utilizing embryos donated from another couple. Eggs from another woman have already been fertilized of the uterus and degree of openness (patency) of the fallopian tubes. transferred to the uterus.

Endometriosis. A condition where endometrial-like tissue (the tissue that lines the uterus) develops outside of the uterine cavity in abnormal locations such as the ovaries, fallopian tubes, and abdominal cavity. This

tissue can grow with hormonal stimulation and cause pain, inflammation, and scar tissue. It may also be associated with infertility.

Estradiol. The predominant estrogen (hormone) produced by the follicles on the ovary.

Fallopian Tubes. A pair of hollow tubes attached one on each side of the uterus through which the egg travels from the ovary to the uterus. Fertilization with the man's sperm usually occurs in the fallopian tube. The fallopian tube is the most common site of ectopic pregnancy.

Fibroids. Benign (non-cancerous) tumors of the uterine muscle wall that can cause abnormal uterine bleeding, pelvic pain, and interfere with pregnancy.

Follicle. A fluid-filled sac located just beneath the surface of the ovary, containing an egg (oocyte) and cells that produce hormones. The sac increases in size and volume during the first half of the menstrual cycle and at ovulation, the follicle matures and ruptures, releasing the egg. As the follicle matures, it can be visualized by ultrasound.

Follicle Stimulating hormone (FSH). In women, FSH is the pituitary hormone responsible for stimulating follicular cells in the ovary to grow, stimulating egg development and the production of the female hormone estrogen. In the male, FSH is the pituitary hormone which travels through the bloodstream to the testes and helps stimulate them to manufacture sperm. FSH can also be given as a medication.

Hysteroscopy. A procedure in which a lighted scope (hysteroscope) is inserted through the cervix into the uterus to enable the physician to view the inside of the uterus to diagnose and treat problems within the uterine cavity.

Hydrosalpinx. A term used to describe a fallopian tube which is swollen or dilated and often filled with fluid. This fluid usually results from a previous infection of the fallopian tube. A hydrosalpinx may be detected by ultrasound, HSG, or by laparoscopy.

Hysterosalpingogram (HSG). An x-ray procedure in which a special media (dye) is injected into the uterus to demonstrate the inner contour of the uterus and degree of openness (patency) of the fallopian tubes.

Intracytoplasmic sperm injection (ICSI). A micromanipulation technique used in conjunction with IVF that involves injecting a sperm directly into an egg in order to facilitate fertilization. The fertilized egg is then transferred to the uterus.

Intrauterine insemination (IUI). The process whereby sperm are injected directly into the uterine cavity in order to bypass the cervix and place the sperm closer to the egg. The sperm are usually washed first in order to remove chemicals that can irritate the uterine lining and to increase sperm motility and concentration.

In vitro fertilization (IVF). A method of assisted reproduction that involves surgically removing eggs from the woman's ovaries, combining them with sperm in the laboratory and, if fertilized, replacing the resulting embryo into the woman's uterus.

Laparoscopy. A narrow, lighted viewing instrument with a telescopic lens.

Laparoscopy. A diagnostic procedure in which a surgeon inserts a laparoscope through a small incision below the navel and visually inspects the uterus, uterine ligaments, fallopian tubes, ovaries, and abdominal organs. Other incisions may also be made to allow insertion of additional instruments to facilitate diagnosis and treatment of pelvic disease.

Luteal phase deficiency. A condition present when the lining of the uterus does not mature properly in response to the progesterone secreted by the ovary after ovulation.

Luteinizing hormone (LH). The hormone that triggers ovulation and stimulates the corpus luteum to secrete progesterone.

Ovary. One of the two female sex glands in the pelvis, located on each side of the uterus. The ovaries produce eggs and hormones including estrogen, progesterone, and androgens.

Ovarian reserve. Refers to a woman's fertility potential in the absence of any problems in the reproductive tract (fallopian tubes, uterus, vagina). It mainly depends on the number and quality of eggs in the ovaries and how well the ovarian follicles are responding to the hormonal signals from the brain.

Ovulation. The release of a ripened egg from its follicle in the outer portion of the ovary. Ovulation usually occurs on day 14 or 15 of a 28-day cycle or 14 days prior to the first day of the next period.

Peritoneum. The lining of the abdominal cavity.

Pre-Implantation Genetic Diagnosis (PGD). A procedure used in conjunction with IVF. At some point 3-5 days after the embryo is formed, part of the embryo is biopsied in order to run special genetic tests. These tests vary and can include counting the number of chromosomes on each embryo or looking specifically for a genetic illness carried by the couple or in the couple's family.

Progesterone. A female hormone secreted by the ovary during the second half of the menstrual cycle after ovulation. Progesterone prepares the lining of the uterus for implantation of the fertilized egg.

Semen analysis. The microscopic examination of semen to determine the number of sperm, their shapes, and their ability to move.

Uterus (womb). The hollow, muscular organ in the pelvis where an embryo implants and grows during pregnancy. The lining of the uterus, called the endometrium, produces the monthly menstrual blood flow when there is no pregnancy.

Vagina. The genital canal in the female that leads from the vulva (external genitalia) to the cervix and uterus.

Varicocele. A varicose or dilated vein within the scrotum that can cause infertility in some men.

For more information on this and other reproductive health topics visit www.ReproductiveFacts.org



Let Us Know What You Think

Email your comments on this booklet to asrm@asrm.org. In the subject line, type “Attention: Patient Education Committee.”

Notes

Notes

Notes



AMERICAN SOCIETY FOR REPRODUCTIVE MEDICINE
1209 MONTGOMERY HIGHWAY
BIRMINGHAM, ALABAMA 35216-2809
(205) 978-5000 • ASRM@ASRM.ORG • WWW.ASRM.ORG